

**AP<sup>®</sup> CALCULUS AB  
2005 SCORING GUIDELINES**


**Question 4**

$x$	0	$0 < x < 1$	1	$1 < x < 2$	2	$2 < x < 3$	3	$3 < x < 4$
$f(x)$	-1	Negative	0	Positive	2	Positive	0	Negative
$f'(x)$	4	Positive	0	Positive	DNE	Negative	-3	Negative
$f''(x)$	-2	Negative	0	Positive	DNE	Negative	0	Positive

Let  $f$  be a function that is continuous on the interval  $[0, 4)$ . The function  $f$  is twice differentiable except at  $x = 2$ . The function  $f$  and its derivatives have the properties indicated in the table above, where DNE indicates that the derivatives of  $f$  do not exist at  $x = 2$ .

(a) For  $0 < x < 4$ , find all values of  $x$  at which  $f$  has a relative extremum. Determine whether  $f$  has a relative maximum or a relative minimum at each of these values. Justify your answer.

(b) On the axes provided, sketch the graph of a function that has all the characteristics of  $f$ .  
(Note: Use the axes provided in the pink test booklet.)




(c) Let  $g$  be the function defined by  $g(x) = \int_1^x f(t) dt$  on the open interval  $(0, 4)$ . For  $0 < x < 4$ , find all values of  $x$  at which  $g$  has a relative extremum. Determine whether  $g$  has a relative maximum or a relative minimum at each of these values. Justify your answer.

(d) For the function  $g$  defined in part (c), find all values of  $x$ , for  $0 < x < 4$ , at which the graph of  $g$  has a point of inflection. Justify your answer.

(a)  $f$  has a relative maximum at  $x = 2$  because  $f'$  changes from positive to negative at  $x = 2$ .

(b)



(c)  $g'(x) = f(x) = 0$  at  $x = 1, 3$ .  
 $g'$  changes from negative to positive at  $x = 1$  so  $g$  has a relative minimum at  $x = 1$ .  $g'$  changes from positive to negative at  $x = 3$  so  $g$  has a relative maximum at  $x = 3$ .

(d) The graph of  $g$  has a point of inflection at  $x = 2$  because  $g'' = f'$  changes sign at  $x = 2$ .

2 : { 1 : relative extremum at  $x = 2$   
1 : relative maximum with justification

2 : { 1 : points at  $x = 0, 1, 2, 3$   
and behavior at  $(2, 2)$   
1 : appropriate increasing/decreasing  
and concavity behavior

3 : { 1 :  $g'(x) = f(x)$   
1 : critical points  
1 : answer with justification

2 : { 1 :  $x = 2$   
1 : answer with justification